

CIRCLE ACTIONS ON TORIC MANIFOLDS AND THEIR APPLICATIONS

Victor M. Buchstaber

Victor.Buchstaber@manchester.ac.uk

Steklov Mathematical Institute,
Russian Academy of Sciences,
Moscow

School of Mathematics,
University of Manchester,
Manchester

Slides for lecture of 17.11.2005, on
joint work with Taras Panov and Nigel Ray.
Mathematics Dept, Leicester University

Prelude to toric manifolds

Consider a smooth $2n$ -dimensional manifold M^{2n} , endowed with a smooth action of the n -dimensional torus T^n .

The orbit space M^{2n}/T^n is a manifold with **corners**.

The most basic manifolds with corners are the **simple convex** polytopes, ie n -dimensional convex polytopes with exactly n facets meeting at each vertex.

A simple convex polytope is **generic**; its bounding hyperplanes are in **general position**.

Each face of a simple polytope is again a simple polytope.

Complex projective space

$$\mathbb{C}P^n = S^{2n+1}/S^1$$

where $S^{2n+1} \subset \mathbb{C}^{n+1}$ consists of vectors

$$z = (z_1, \dots, z_{n+1}) \quad \text{with} \quad |z|^2 = \sum_{k=1}^{n+1} z_k^2 = 1,$$

and $t_1 \in S^1$ acts by $t_1 \cdot z = (t_1 z_1, \dots, t_1 z_{n+1})$.

The **standard action** of T^{n+1} on \mathbb{C}^{n+1} is

$$t \cdot z = (t_1 z_1, \dots, t_{n+1} z_{n+1}),$$

and induces an action of T^n on $\mathbb{C}P^n$ by

$$t \cdot [z] = [t_1 z_1, \dots, t_n z_n, z_{n+1}].$$

The orbit space is the n -**simplex** Δ^n , given by

$$\{(x_1, \dots, x_{n+1}) \in \mathbb{R}^{n+1}, x_i \geq 0, \sum_{k=1}^{n+1} x_k = 1\}.$$

The projection $\pi: \mathbb{C}P^n \rightarrow \Delta^n$ acts by

$$(z_1 : z_2 : \dots : z_{n+1}) \longrightarrow (|z_1|^2, |z_2|^2, \dots, |z_{n+1}|^2)$$

The stable tangent bundle of $\mathbb{C}P^n$

We write the Hopf bundle as

$$p : \eta \longrightarrow \mathbb{C}P^n.$$

A point of η is a pair (L, v) , where L is a line in \mathbb{C}^{n+1} and v is a point in L . The projection acts by $p(L, v) = L$. There is an isomorphism

$$\tau(\mathbb{C}P^n) \oplus \mathbb{C} \simeq (n + 1)\bar{\eta},$$

where $\bar{\eta} = \text{Hom}(\eta, \mathbb{C})$.

The bounded flag manifolds B_n

A point

$$U = (U_1 \subset U_2 \subset \cdots \subset U_{n+1}) \in B_n$$

is a **complete flag** in \mathbb{C}^{n+1} such that each U_k contains the subspace \mathbb{C}^{k-1} , spanned by the first $k - 1$ standard basis vectors in \mathbb{C}^{n+1} , for $2 \leq k \leq n$. So U is equivalent to a sequence of lines

$$\{L_k \subset \mathbb{C}_k \oplus L_{k+1}, 1 \leq k \leq n\}$$

where \mathbb{C}_k is k -th coordinate line and $L_1 = U_1$, $L_{n+1} = \mathbb{C}_{n+1}$. The standard action of T^{n+1} on \mathbb{C}^{n+1} induces an action of T^n on B_n , whose orbit space is the **cube** $I^n = I \times \cdots \times I$.

The projection $\pi: B_n \rightarrow I^n$ satisfies

$$\pi(U) = (\pi(L_1), \dots, \pi(L_n)),$$

where $\pi(L_k) = |l_k|^2$ and l_k is the projection of a unit vector from $L_k \subset \mathbb{C}_k \oplus L_{k+1}$ into \mathbb{C}_k .

The stable tangent bundle of B_n

For each $1 \leq k \leq n$, a complex line bundle

$$p_\gamma: \gamma_k(n) \longrightarrow B_n$$

is defined by $p_\gamma^{-1}(U) = (U, L_k)$.

We set $\gamma_{n+1}(n)$ to be trivial.

Each B_n is the 2-sphere bundle of

$$\gamma_1(n) \oplus \mathbb{R} \longrightarrow B_{n-1}.$$

By Szczarba's construction, this leads to a **stably complex structure**

$$\tau(B_n) \oplus \mathbb{R}^2 \simeq \bigoplus_{k=2}^{n+1} \gamma_k(n) \oplus \mathbb{C},$$

which extend over the disc bundle of $\gamma_1(n) \oplus \mathbb{R}$.

The manifolds $B_{i,j}$

$B_{i,j}$ is a $2n$ -dimensional smooth $\mathbb{C}P^{j-1}$ -bundle over B_i , where $n = (i + j - 1)$ and $0 \leq i \leq j$.

Each point of $B_{i,j}$ is a pair (U, W) , where $U \in B_i$ is a bounded flag in \mathbb{C}^{i+1} and W is a line in $U_1^\perp \oplus \mathbb{C}^{j-i}$. The projection

$$\pi: B_{i,j} \longrightarrow B_i$$

satisfies $\pi(U, W) = U$.

So $T^n = T^i \times T^{j-1}$ acts on $B_{i,j}$ by

$$(t_1 z_1, \dots, t_i z_i, z_{i+1}, t_{i+1} w_1, \dots, t_n w_{j-1}, w_j).$$

The orbit space is $I^i \times \Delta^{j-1}$, and the quotient map

$$B_{i,j} \longrightarrow I^i \times \Delta^{j-1}$$

is given by $\pi(U, W) = (\pi(U), \pi(W))$.

The stable tangent bundle of $B_{i,j}$

We define a complex line bundle ζ over $B_{i,j}$ by considering W as a line in \mathbb{C}^{j+1} . The projection $\pi: B_{i,j} \rightarrow B_i$ and the classifying map $B_{i,j} \rightarrow \mathbb{C}P^j$ of ζ together provide a smooth embedding

$$B_{i,j} \hookrightarrow B_i \times \mathbb{C}P^j,$$

whose normal bundle is the pullback of $\gamma_1(i) \otimes \zeta$.

The resulting isomorphism

$$\tau(B_{i,j}) \oplus (\gamma_1(i) \otimes \zeta) \oplus \mathbb{C} \simeq \bigoplus_{k=2}^{i+1} \gamma_k(i) \oplus (j+1)\bar{\zeta}$$

defines a **stably complex structure** on $B_{i,j}$.

Moment-angle manifolds

We deal only with simple polytopes, and reserve the notation $m = m(P)$ for the number facets of P . Every face of codimension k may be written uniquely as

$$F_I = F_{i_1} \cap \cdots \cap F_{i_k} \quad (1)$$

for some subset $I = \{i_1, \dots, i_k\}$ of $[m]$.

We denote the i -th coordinate subcircle of the standard m -torus T^m by T_i for every $1 \leq i \leq m$. Given any subset $I \subseteq [m]$, we define the subgroup by

$$T_I = \prod_{i \in I} T_i < T^m$$

in particular, T_\emptyset is the trivial subgroup $\{1\}$.

Every point p of P lies in the interior of a unique face F_{I_p} , where $I_p = \{i : p \in F_i\}$, and it is convenient to abbreviate F_{I_p} to $F(p)$ and T_{I_p} to $T(p)$. If p is a vertex, then $T(p)$ has dimension n (the maximum possible), and if p is an interior point of P , then $T(p)$ is trivial.

We now construct the identification space

$$\mathcal{Z}_P = T^m \times P / \sim \quad (2)$$

where $(t_1, p) \sim (t_2, p)$ if and only if

$$t_1^{-1}t_2 \in T(p).$$

So \mathcal{Z}_P is an $(m + n)$ -dimensional manifold with a canonical left T^m -action, whose isotropy subgroups are precisely the subgroups $T(p)$.

Dicharacteristics

In order to construct toric manifolds over P , we need one further set of data. This consists of a homomorphism $\ell: T^m \rightarrow T^n$, whose properties are controlled by P (cf. Davis and Januszkiewicz), namely

$$F_I \text{ a face of codim } k \Rightarrow \ell \text{ monic on } T_I. \quad (3)$$

Any such ℓ is called a **dicharacteristic**; condition (3) ensures that the kernel $K(\ell)$ of ℓ is isomorphic to an $(m - n)$ -dimensional subtorus of T^m . Wherever possible we abbreviate $K(\ell)$ to K .

We write the subcircle $\ell(T_i) < T^n$ as $T(F_i)$ for any $1 \leq i \leq m$, and the subgroup $\ell(T_I)$ as $T(F_I)$ for any face F_I . For each point p in P let $S(p)$ denote the subgroup $T(F(p))$; it is, of course, $\ell(T(p))$. For example, $S(w) = T^n$ for any vertex w , and $S(p) = \{1\}$ for any point p in the interior of P .

Refined form of dicharacteristic matrix

Applied to the initial vertex $v_\star = F_1 \cap \cdots \cap F_n$, (3) ensures that the restriction of ℓ to $T_1 \times \cdots \times T_n$ is an isomorphism. So we may use the circles $T(F_1), \dots, T(F_n)$ to define a basis for the Lie algebra of T^n , and represent the homomorphism induced by ℓ by an $n \times m$ integral matrix

$$\Lambda = \begin{pmatrix} 1 & 0 & \cdots & 0 & \lambda_{1,n+1} & \cdots & \lambda_{1,m} \\ 0 & 1 & \cdots & 0 & \lambda_{2,n+1} & \cdots & \lambda_{2,m} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 & \lambda_{n,n+1} & \cdots & \lambda_{n,m} \end{pmatrix}. \quad (4)$$

Given any other vertex $v = F_{j_1} \cap \cdots \cap F_{j_n}$, (3) implies that the corresponding columns $\lambda_{j_1}, \dots, \lambda_{j_n}$ form a basis for \mathbb{Z}^n , and have determinant ± 1 . We refer to (4) as the **refined form** of the dicharacteristic matrix.

A model for toric manifolds

Since $K = K(\ell)$ acts freely on \mathcal{Z}_P there is a principal K -bundle

$$\pi_\ell: \mathcal{Z}_P \rightarrow M,$$

whose base space is a $2n$ -dimensional manifold. By construction,

$$M = T^n \times P / \sim \tag{5}$$

where $(s_1, p) \sim (s_2, p)$ if and only if

$$s_1^{-1} s_2 \in S(p).$$

Furthermore, M admits a canonical T^n -action α , which is locally isomorphic to the standard action on \mathbb{C}^n , and has quotient map

$$\pi: M \rightarrow P.$$

Note that $\pi \cdot \pi_\ell$ is the natural projection $\varrho_P: \mathcal{Z}_P \rightarrow P$.

The fixed points of α project to the vertices of P , and we refer to $\pi^{-1}(v_\star)$ as the **initial fixed point** x_\star .

Then (5) identifies a neighbourhood of x_\star with \mathbb{C}^n , on which α is standard; its representation at other fixed points $\pi^{-1}(v)$ may be read off from the corresponding columns of Λ .

The quadruple (M, α, π, P) is an example of Davis and Januszkiewicz's **toric manifolds**.

Any manifold with a similarly well-behaved torus action over P is equivariantly diffeomorphic to a model of the form (5).

Facial submanifold structure

The **facial submanifolds** M_i of M are defined as the inverse images of the facet F_i under π , for $1 \leq i \leq m$. Every M_i has codimension 2, with isotropy subgroup $T(F_i) < T^n$. The quotient map

$$\mathcal{Z}_P \times_K \mathbb{C}_i \longrightarrow M \tag{6}$$

defines a canonical complex line-bundle ρ_i , whose restriction to M_i is isomorphic to the normal bundle ν_i of its embedding in M .

The submanifolds M_i are mutually transverse, and we write any non-empty intersection as

$$M_I = M_{i_1} \cap \cdots \cap M_{i_k}. \tag{7}$$

So M_I is the inverse image under π of the codimension- k face F_I . M_I has codimension $2k$, and its isotropy subgroup is $T(F_I)$.

The restriction of $\rho_I = \rho_{i_1} \oplus \cdots \oplus \rho_{i_d}$ to M_I is isomorphic to the normal bundle ν_I of its embedding in M , for any face F_I .

The cohomology ring of a toric manifold

The bundles ρ_i are important in understanding the integral cohomology ring of M .

Let u_i be the first Chern class $c_1(\rho_i)$ in $H^2(M)$; then $H^*(M)$ is generated by the elements u_1, \dots, u_m , modulo two sets of relations.

The first are linear, and arise from the refined form (4) of the dicharacteristic; the second are monomial, and arise from the Stanley-Reisner ideal of P .

The linear relations take the form

$$u_i = -\lambda_{i,n+1}u_{n+1} - \dots - \lambda_{i,m}u_m, \quad (8)$$

for $1 \leq i \leq n$. So u_{n+1}, \dots, u_m suffice to generate $H^*(M)$ multiplicatively.

Fixed points of subcircle actions

Given the action α of T^n on M , it is natural to study its restriction to an arbitrary subcircle $T < T^n$. We may decompose the **fixed point set** of T into the union of its components as

$$\text{Fix}(T) = M_{I(1)} \cup \cdots \cup M_{I(s)} \cup \cdots \cup M_{I(d)}, \quad (9)$$

Following (4), we represent T by a primitive column vector $l = l(T)$ in \mathbb{Z}^n .

Proposition. The components of $\text{Fix}(T)$ are specified by those $I(s)$ for which none of the coefficients $\alpha_{i(s)j}$ is zero in any expansion of the form

$$l(T) = \alpha_{i(s)_1} \lambda_{i(s)_1} + \cdots + \alpha_{i(s)_k} \lambda_{i(s)_k}, \quad (10)$$

for $1 \leq s \leq d$.

Stably complex, special unitary, and level- L structures

On a smooth manifold N of dimension d , a **stably complex structure** is an equivalence class of real $2k$ -plane bundle isomorphisms

$$\tau(N) \oplus \mathbb{R}^{2k-d} \simeq \zeta,$$

where ζ is a fixed $GL(k, \mathbb{C})$ -bundle over N and k is suitably large. Two such isomorphisms are equivalent when they agree up to stabilisation.

If the first Chern class $c_1(\zeta)$ is zero, then the stably complex structure is **special unitary**, (or SU); and if it is divisible by a positive integer L , then it is **level- L** .

We identify the geometric data required to induce such structures on a toric manifold.

Note that $\mathbb{C}P^n$ is level- L for any divisor L of $n + 1$, where $n \geq 1$.

Combinatorial data underlying an omnioriented toric manifold

An **omniorientation** of a toric manifold M consists of a choice of orientation for M , and for every normal bundle ν_i .

An interior point of the quotient polytope P admits an open neighborhood U , whose inverse image under the projection π is canonically diffeomorphic to $T^n \times U$ as a subspace of M . Since T^n is oriented by the standard choice of basis, orientations of M correspond bijectively to orientations of P . Moreover, the dicharacteristic ℓ determines a complex structure on every ρ_i , so it encodes an orientation for every ν_i .

Every pair (P, Λ) therefore determines a $2n$ -dimensional **omnioriented toric manifold**, where P is the combinatorial type of an oriented finely ordered n -dimensional simple polytope, and Λ is a matrix of the form (4).

Stably complex structures

Theorem. Any omnioriented toric manifold admits a canonical stably complex structure, which is invariant under the T^n -action.

Proof. Using the theory of **analogous polytopes** we obtained an embedding

$$i_P: P \longrightarrow \mathbb{R}_{\geq}^m$$

which respects facial codimensions and gives a pullback diagram

$$\begin{array}{ccc} \mathcal{Z}_P & \xrightarrow{i_Z} & \mathbb{C}^m \\ \varrho_P \downarrow & & \downarrow \varrho \\ P & \xrightarrow{i_P} & \mathbb{R}^m \end{array} \quad (11)$$

of identification spaces. Here $\varrho(z_1, \dots, z_m)$ is given by $(|z_1|^2, \dots, |z_m|^2)$, the vertical maps are projections onto the quotients by the T^m -actions, and i_Z is a T^m -equivariant embedding.

So, there is a K -**equivariant** decomposition

$$\tau(\mathcal{Z}_P) \oplus \nu(i_Z) \simeq \mathcal{Z}_P \times \mathbb{C}^m,$$

obtained by restricting the tangent bundle $\tau(\mathbb{C}^m)$ to \mathcal{Z}_P . Factoring out K yields

$$\tau(M) \oplus (\xi/K) \oplus (\nu(i_Z)/K) \simeq \mathcal{Z}_P \times_K \mathbb{C}^m, \quad (12)$$

where ξ denotes the $(m - n)$ -plane bundle of tangents along the fibres of

$$\pi_\ell : \mathcal{Z}_P \longrightarrow M.$$

The right-hand side of (12) is isomorphic to $\bigoplus_{i=1}^m \rho_i$ as $GL(m, \mathbb{C})$ -bundles. This is all an example of Szczarba's Theorem.

The embedding $i_{\mathbb{Z}} : \mathcal{L}_P \longrightarrow \mathbb{C}^m \simeq \mathbb{R}^{2n}$ is T^m -equivariantly framed, so $\nu(i_{\mathbb{Z}})/K$ is trivial. The bundle ξ/K canonically isomorphic to the adjoint bundle of the principal K -bundle, which is trivial because K is abelian.

So, (12) reduces to an isomorphism

$$\tau(M) \oplus \mathbb{R}^{2(m-n)} \simeq \rho_1 \oplus \dots \oplus \rho_m,$$

although different choices of trivialisations may lead to different isomorphisms.

Since M is connected and $GL(2(m-n), \mathbb{R})$ has two connected components, such isomorphisms are equivalent when and only when the induced orientations agree on $\mathbb{R}^{2(m-n)}$.

We choose the orientation which is compatible with those on $\tau(M)$ and $\rho_1 \oplus \dots \oplus \rho_m$, as given by the omniorientation. The induced structure is invariant under the action of T^n , because i_Z is T^m -equivariant.

SU, and level- L structures

Corollary. The omniorientation induces an SU -structure on M precisely when the refined matrix Λ of (4) has every column-sum equal to 1; it induces a level- L structure when every column-sum is congruent to 1 mod L .

Proof. The stably complex structure induced by the omniorientation has first Chern class $\sum_{i=1}^m u_i$. It is zero in $H^2(M)$ if and only if

$$\left(1 - \sum_{i=1}^n \lambda_{i,n+1}\right) u_{n+1} + \cdots + \left(1 - \sum_{i=1}^n \lambda_{i,m}\right) u_m = 0.$$

The same argument shows that it is divisible by L if and only if every column-sum is congruent to 1 mod L .

Applications to complex cobordism

Theorem. Every complex cobordism class is represented by a disjoint union of omnioriented toric manifolds, which are suitably oriented products of the $B_{i,j}$.

Our modification of **connected sum** of toric manifolds gives:

Theorem. In dimensions > 2 , every complex cobordism class contains a toric manifold, necessarily connected, whose stably complex structure is induced by an omniorientation, and is therefore compatible with the action of the torus.

Similar methods combine with Diagram (11) to give:

Theorem. Every complex cobordism class may be represented by the quotient of a free torus action on a real quadratic complete intersection.

Formal groups

Let

$$\mathcal{F}(u, v) = u + v + \sum a_{ij} u^i v^j \in \Omega_U[[u, v]],$$

with $\deg u = \deg v = 2$ and $\deg a_{ij} = -2(i + j - 1)$, be the **formal group** of complex cobordism theory, and

$$g(u) = u + \sum [CP(n)] \frac{u^{n+1}}{n+1} \in \Omega_U \otimes Q[[u]]$$

its **logarithm**. So

$$g'(u) \in \Omega_U[[u]] \quad \text{and} \quad \mathcal{F}'_v(u, 0) = \frac{1}{g'(u)}.$$

The **universality** of $\mathcal{F}(u, v)$ implies that, for every formal group law $f(u, v)$ over a ring R , there exists a ring homomorphism

$$\Omega_U \longrightarrow R$$

classifying $f(u, v)$. Such a homomorphism is otherwise known as a **Hirzebruch genus**.

Hirzebruch genera

Every Hirzebruch genus $T_h: \Omega_U \longrightarrow R$ may also be defined in terms of a series $x/h(x)$, where

$$h(x) = x + \sum_{i=1}^{\infty} \lambda_i x^{i+1} \in R \otimes \mathbb{Q}[[x]].$$

If T_h classifies the formal group law $f(u, v)$, then $h(x)$ is the compositional inverse to the logarithm $h(x)$, and is its **exponential series**.

Then T_h is evaluated on stably complex manifolds M^{2n} by the formula

$$T_h(M^{2n}) = \left\langle \prod_{j=1}^n \frac{x_j}{h(x_j)}, [M^{2n}] \right\rangle,$$

where $\prod_{j=1}^n (1 + x_j) = 1 + \sum c_k(M^{2n})$, and the $c_k(M^{2n})$ are the **Chern classes** of the stable complex tangent bundle of M^{2n} .

Important examples

1. Consider the formal group law

$$f_{\alpha,\beta}(u, v) = \frac{u + v - auv}{1 - buv},$$

with $a = \alpha + \beta$, $b = \alpha\beta$, $\deg a = -2$, and $\deg b = -4$. Then

$$g'_f(u) = \frac{1}{\alpha - \beta} \left(\frac{\alpha}{1 - \alpha u} - \frac{\beta}{1 - \beta u} \right) = 1 + \sum T_{\alpha,\beta}[CP^n]$$

where $T_{\alpha,\beta}$ is the **two-parameter Todd genus**.

In this case, the exponential series is given by

$$h(x) = \frac{\alpha \exp \frac{\alpha - \beta}{2} x - \beta \exp \frac{\beta - \alpha}{2} x}{\exp \frac{\alpha - \beta}{2} x - \exp \frac{\beta - \alpha}{2} x}.$$

2. Consider the formal group law

$$f_{ell}(u, v) = \frac{u\sqrt{R(v)} + v\sqrt{R(u)}}{1 - \varepsilon u^2 v^2},$$

with $R(u) = 1 - 2\delta u^2 + \varepsilon u^4$ and $\deg \delta = -4$,
 $\deg \varepsilon = -8$. Then

$$g'_{ell}(u) = \frac{1}{\sqrt{R(u)}} = 1 + \sum T_{ell}[CP^n]u^n,$$

where T_{ell} is the **elliptic genus**.

In this case, the exponential series

$$h(x) = \operatorname{sn}(x)$$

is the **Jacobi elliptic function** sn .

The generalized elliptic genus

Krichever's generalized elliptic genus T_{ell*} is a Hirzebruch genus T_h , with $h(x) = \frac{1}{\Phi(x)}$ and

$$\Phi(x) = \Phi(x, z) = \frac{\sigma(z-x)}{\sigma(x)\sigma(z)} \exp \zeta(z)x.$$

Here $\sigma(z)$ is the classical **Weierstrass function** and $\zeta(z) = (\ln \sigma(z))'$.

The function $\Phi(x, z)$ is a solution of the **Lame equation**

$$\left(\frac{d^2}{dx^2} - 2\wp(x) \right) \Phi(x) = \wp(z)\Phi(x)$$

of the form

$$\Phi(x) = \frac{1}{x} + \Phi_{reg}(x)$$

in a neighbourhood of $x = 0$, where $\Phi_{reg}(x)$ is a power series such that $\Phi_{reg}(0) = 0$. As usual, $\wp(z) = -\zeta(z)'$.

The power series of the sigma-function

The function $\sigma(x)$ is associated to the universal elliptic curve $\gamma^2 = 4\xi^3 - g_2\xi - g_3$, and according to K. Weierstrass, Zür Theorie der elliptischen Funktionen, *Mathematische Werke*, (Teubner, Berlin, 1894), Vol 2, 245-255, may be expanded as

$$\sigma(x) = x \sum_{i,j \geq 0} \frac{n_{i,j}}{(4i + 6j + 1)!} \left(\frac{g_2 x^4}{2} \right)^i (2g_3 x^6)^j.$$

The $n_{i,j}$ are integers, defined recursively by

$$\begin{aligned} n_{i,j} = & 3(i + 1)n_{i+1,j-1} + \frac{16}{3}(j + 1)n_{i-2,j+1} \\ & - \frac{1}{3}(2i + 3j - 1)(4i + 6j - 1)n_{i-1,j} \end{aligned}$$

plus the initial conditions $n_{0,0} = 1$ and $n_{i,j} = 0$ for $\min(i, j) < 0$.

So we may interpret

$$\sigma(x) = \sigma(x, g_2, g_3) \in \mathbb{Q}[g_2, g_3][[x]]$$

as a homogeneous series of degree 2, with $\deg x = 2$, $\deg g_2 = -8$, and $\deg g_3 = -12$.

The power series of the Krichever genus

Put

$$h(x) = \sigma(x) \exp \varphi(z, x),$$

where

$$\exp \varphi(z, x) = \frac{\sigma(z)}{\sigma(z-x)} \exp(-\zeta(z)x).$$

Then

$$\begin{aligned} \varphi(z, x) &= \ln \sigma(z) - \ln \sigma(z-x) - \zeta(z)x \\ &= \sum_{k=2}^{\infty} (-1)^k \left(-\frac{d^k \ln \sigma(z)}{dz} \right) \frac{x^k}{k!} \\ &= \sum_{k=1}^{\infty} (-1)^{k-1} \wp^{(k-1)}(z) \frac{x^{k+1}}{(k+1)!}. \end{aligned}$$

Writing $\wp(z) = \xi$ and $\wp'(z) = \gamma$, we will show that

$$\varphi(z, x) \in \mathbb{Q}[\xi, \gamma, g_2][[x]]$$

is a homogeneous series of degree 0, with $\deg x = \deg z = 2$, $\deg \xi = -4$, $\deg \gamma = -6$, and $\deg g_2 = -8$.

Using the Weierstrass uniformization

$$(\wp'(z))^2 = 4\wp(z)^3 - g_2\wp(z) - g_3,$$

we obtain

$$\wp''(z) = 6\wp(z)^2 - g_2/2.$$

Then

$$\wp^{(k)}(z) = p_k(\xi, \gamma, g_2) \in \mathbb{Q}[\xi, \gamma, g_2],$$

where $p_1(\xi, \gamma, g_2) = \xi$ and

$$p_{k+1} = \gamma \frac{\partial}{\partial \xi} p_k + \left(6\xi^2 - \frac{g_2}{2}\right) \frac{\partial}{\partial \gamma} p_k$$

for $k \geq 1$. So

$$\varphi(z, x) = \sum_{k=1}^{\infty} (-1)^{k-1} p_k(\xi, \gamma, g_2) \frac{x^{k+1}}{(k+1)!}.$$

Using $g_3 = 4\xi^3 - \gamma^2 - g_2\xi$ we deduce that $\sigma(x) \in \mathbb{Q}[\xi, \gamma, g_2][[x]]$.

Corollary. The Krichever genus T_{ell^*} is a Hirzebruch genus T_h , with

$$h(x) \in \mathbb{Q}[\xi, \gamma, g_2][[x]].$$

The formal group of the generalized elliptic genus

Over the graded ring $\mathbf{R} = \sum_{i \geq 0} \mathbf{R}^{-2i}$, let

$$c_i(u) = \sum_{j \geq 0} c_{i,j} u^j, \quad i = 1, 2,$$

where $\deg c_{1,j} = -2j$, $\deg c_{2,j} = -2j - 4$ and $c_{1,0} = 1$, $c_{1,1} = 0$, $c_{2,0} = 0$.

Theorem. (cf. V.M.Buchstaber, Russian Math. Surveys, 45:3, 1990, 213-215.) The universal formal group $f(u, v)$ of the form

$$uc_1(v) + vc_1(u) - auv - \frac{c_2(u) - c_2(v)}{uc_1(v) - vc_1(u)} u^2 v^2$$

over $\mathbf{R} = \mathbb{Z}[a, c_{1,j}, j \geq 2, c_{2,k}, k \geq 1]/J$ has exponential series

$$h(x) = \frac{\sigma(x)\sigma(z)}{\sigma(z-x)} \exp(a - \zeta(z)) \in \mathbb{Q}[a, \xi, \gamma, g_2][[x]],$$

where J is determined by the associativity conditions, and

$$\mathbf{R} \otimes \mathbb{Q} = \mathbb{Q}[a, c_{1,2}, c_{1,3}, c_{1,4}].$$

For the Krichever genus

$$T_h: \Omega_U \longrightarrow \mathbf{R} \quad \text{with} \quad h(x) = \frac{\exp ax}{\Phi(z, x)},$$

we have

Corollary.

$$1 + \sum_{j \geq 1} T_h(a_{j,1})u^j = c_1(u) - au$$

and

$$\sum_{j \geq 1} T_h(a_{j,2})u^{j-1} = c_{1,2} - c_2(u).$$

Corollary. The homomorphism

$$T_h: \sum_{i=0}^4 \Omega_U^{-2i} \longrightarrow \sum_{i=0}^4 \mathbf{R}^{-2i}$$

is an isomorphism.

Important examples:

1. The formal group law

$$f_{\alpha,\beta}(u, v) = \frac{u + v - auv}{1 - buv}$$

is given by

$$c_1(u) = 1 - bu^2 \quad \text{and} \quad c_2(u) = -bu(a + bu).$$

2. The formal group law

$$f(u, v) = \frac{u\sqrt{R(v)} + v\sqrt{R(u)}}{1 - \varepsilon u^2 v^2},$$

with $R(u) = 1 - 2\delta u^2 + \varepsilon u^4$, is given by

$$c_1(u) = \sqrt{R(u)}, \quad c_2(u) = -\varepsilon u^2, \quad \text{and} \quad a = 0.$$

Krichever's results

Let F_s be a connected component of the set $\{F_s\}$ of fixed points under the action of S^1 on a stably complex S^1 -manifold M .

Suppose that the representation of S^1 in the normal bundle to F_s is given by $\sum_i \eta^{j_{s,i}}$. Then if $c_1(M) = 0$, all the sums $r_s = \sum_i j_{s,i}$ are equal. The resulting integer is called the **type** of the circle action on the SU -manifold M .

Theorem. If the action of S^1 on any SU -manifold M has nonzero type, then

$$T_{ell_*}([M]) = 0,$$

where T_{ell_*} is Krichever's generalized elliptic genus.

The generalized elliptic genus of a toric manifold

Let M^{2n} be a toric SU -manifold and $T < T^n$ be an arbitrary subcircle. So

$$l(T) = (z_1, \dots, z_n) \in \mathbb{Z}^n.$$

Proposition. The action of T on M^{2n} has type $\sum_{i=1}^n z_i$.

Corollary. The generalized elliptic genus T_{ell*} of any toric SU -manifold M^{2n} is zero.

Corollary. If M^{2n} is any toric SU -manifold, where $n < 5$, then $[M^{2n}] = 0$.

References

1. Victor M Buchstaber, Taras E Panov and Nigel Ray. Analogous Polytopes, Circle Actions, and Toric Manifolds. *in preparation*.
2. Victor M Buchstaber and Taras E Panov. Torus Actions and Their Applications in Topology and Combinatorics. Volume 24 of *University Lecture Series*, Amer Math Soc, Providence, RI, 2002.
3. Victor M Buchstaber and Nigel Ray. Tangential structures on toric manifolds, and connected sums of polytopes. *Internat Math Res Notices*, 4:193–219, 2001.
4. Victor M Buchstaber and Nigel Ray. Toric Manifolds and Complex Cobordism (Russian). *Uspekhi Mat Nauk* 53(2):139–140, 1998. (English translation in *Russian Math Surveys* 53(2):371–373, 1998.)